

Enterisol® Ileitis

Enterisol® Ileitis FF

Lawsonia Intracellularis Vaccine, Avirulent Live Culture

Description

Enterisol® Ileitis is a safe, effective, convenient, one dose, live *Lawsonia intracellularis* (ileitis) vaccine. It is administered orally.

Features

- Safe for pigs 3 weeks of age or older.
- Single dose administration.
- Easy to use, oral administration via the drinking water.
- Can be administered via oral drench.
- Avirulent live culture vaccine.

Benefits

- Significantly reduces gross and microscopic intestinal lesions of ileitis.
- Significantly reduces colonization by *Lawsonia intracellularis* following challenge with virulent microorganism.

Advantages

- Orally administered, modified-live ileitis vaccine.
- Convenient water delivery results in less animal stress.
- Simple oral vaccination reduces the need for injections.
- Less exposure to needles results in improved pork quality and worker safety.

Dosage & Administration

Administer one dose via drinking water or oral drench to each pig.

Lyophilized Form – Rehydrate the vaccine by adding the full contents of the accompanying sterile diluent to the vaccine. Shake well and use entire contents immediately. The Lyophilized form may be administered via oral drench or water.

Frozen Form (FF) – Thaw vaccine under trickling cool water bath until liquid (usually 30-40 minutes). Shake well and use entire contents immediately.

See product insert for complete directions.

Precautions

Lyophilized Form – Store at 35°– 46°F. Do not freeze. Use entire contents immediately after reconstituting.

Frozen Form – Store frozen at ≤ -70°C. Do not thaw until immediately prior to use.

Use entire contents immediately after thawing. Do not refreeze.

In case of anaphylactoid reaction, epinephrine is symptomatic treatment.

Highlights

- Avirulent live culture vaccine
- Drinking water administration
- Oral drench administration capability
- Less animal stress during vaccination
- PQA friendly - no needles

Uses

Enterisol® Ileitis is recommended for the vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the prevention and control of porcine proliferative enteropathy (Ileitis) caused by *Lawsonia intracellularis*.

Can be used in conjunction with Reload Pack®.

Warning

For veterinary use only.

Do not vaccinate within 21 days of slaughter.


This product has not been tested in pregnant animals.

Do not use concurrently with antibiotics.

Product Information

Name	Code	Size
ENTERISOL ILEITIS	137724	100mL 50 dose
ENTERISOL ILEITIS	137949	100mL 100 dose
ENTERISOL ILEITIS FF	137188	100mL 100 dose
ENTERISOL ILEITIS FF	137438	100mL 250 dose
ENTERISOL ILEITIS FF	136476	100mL 500 dose
RELOAD PACK Blue Dye/Chlorine Neutralizer	136904	1 gallon
RELOAD PACK Blue Dye/Chlorine Neutralizer	138702	5 gallon



<p>Serial No.: Exp. Date:</p>	<p>This product has been shown to be effective for the vaccination of healthy swine 3 weeks of age or older against <i>Lawsonia intracellularis</i>. The duration of immunity has not been determined. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.</p> <p>The vaccine has been shown to have an effect on gross and microscopic intestinal lesions of ileitis.</p> <p>Directions and dosage: Rehydrate the vaccine by adding the accompanying diluent to the vaccine vial. Mix well and use immediately. Administer one 2 mL dose per pig, either orally or via the drinking water. See inside of carton for complete directions.</p> <p>Precautions: Store at 35-46°F (2-8°C). Do not freeze. Use entire contents when first opened.</p>		<p>Do not vaccinate within 21 days before slaughter. Do not mix with other products, except as specified on this label. Do not use in pregnant swine or breeding boars. Do not use concurrently with antimicrobials effective against <i>Lawsonia</i> spp. All materials used in administering this vaccine must be free of such antimicrobial or disinfectant residue to prevent inactivation and reduced product efficacy. In case of anaphylactoid reaction, administer epinephrine. In case of human exposure, contact a physician. Inactivate unused contents before disposal.</p> <p>Boehringer Ingelheim Animal Health USA Inc. St. Joseph, MO 64506 Phone: 1 (888) 637-4251 VLN/PCN 124/10L1.01</p> 	130402-03
-----------------------------------	---	---	---	-----------

<p>Serial No.: Exp. Date:</p>	<p>This product has been shown to be effective for the vaccination of healthy swine 3 weeks of age or older against <i>Lawsonia intracellularis</i>. The duration of immunity has not been determined. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.</p> <p>The vaccine has been shown to have an effect on gross and microscopic intestinal lesions of ileitis.</p> <p>Directions and dosage: Rehydrate the vaccine by adding the accompanying diluent to the vaccine vial. Mix well and use immediately. Administer one 1 mL dose per pig, either orally or via the drinking water. See inside of carton for complete directions.</p> <p>Precautions: Store at 35-46°F (2-8°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter.</p>		<p>Do not mix with other products, except as specified on this label. Do not use in pregnant swine or breeding boars. Do not use concurrently with antimicrobials effective against <i>Lawsonia</i> spp. All materials used in administering this vaccine must be free of such antimicrobial or disinfectant residue to prevent inactivation and reduced product efficacy. In case of anaphylactoid reaction, administer epinephrine. In case of human exposure, contact a physician. Inactivate unused contents before disposal.</p> <p>Boehringer Ingelheim Animal Health USA Inc. St. Joseph, MO 64506 Phone: 1 (888) 637-4251 VLN/PCN 124/10L1.01</p> 	130403-04
-----------------------------------	---	---	---	-----------

<p>Serial No.: Exp. Ref./Batch:</p>	<p>0 12313129611 3</p> 		<p>For use in animals only. Store frozen at ≤ -60°C. Do not vaccinate within 21 days before slaughter. Inactivate unused contents before disposal. See insert for complete information.</p> <p>Pour utilisation chez les animaux seulement. Entreposer au congélateur à une température de -60 °C ou moins. Ne pas vacciner dans les 21 jours précédant l'abattage. Inactiver tout contenu inutilisé avant son élimination. Voir la notice pour l'information complète.</p> <p>Ver las instrucciones completas en el Instructivo adjunto.</p> <p>Manufactured by/Fabriqué par/Fabricado por: Boehringer Ingelheim Animal Health USA Inc. St. Joseph, MO 64506 Tel/Fel.: 1 (888) 637-4251 VLN/PCN 124/10L1.00</p> <p>Distributed in Canada by/Distribué au Canada par/Distribuido en Canada por: Boehringer Ingelheim Animal Health Canada Inc. Boehringer Ingelheim Santé Animale Canada Inc. Burlington ON L7L 5H4</p> 	129621-03
---	---	--	---	-----------

<p>Serial No.: Exp. Ref./Batch:</p>	<p>0 12313129620</p> 		<p>For use in animals only. Store frozen at ≤ -60°C. Do not vaccinate within 21 days before slaughter. Inactivate unused contents before disposal. See insert for complete information.</p> <p>Pour utilisation chez les animaux seulement. Entreposer au congélateur à une température de -60 °C ou moins. Ne pas vacciner dans les 21 jours précédant l'abattage. Inactiver tout contenu inutilisé avant son élimination. Voir la notice pour l'information complète.</p> <p>Ver las instrucciones completas en el Instructivo adjunto.</p> <p>Manufactured by/Fabriqué par/Fabricado por: Boehringer Ingelheim Animal Health USA Inc. St. Joseph, MO 64506 Tel/Fel.: 1 (888) 637-4251 VLN/PCN 124/10L1.00</p> <p>Distributed in Canada by/Distribué au Canada par/Distribuido en Canada por: Boehringer Ingelheim Animal Health Canada Inc. Boehringer Ingelheim Santé Animale Canada Inc. Burlington ON L7L 5H4</p> 	129622-03
---	--	---	---	-----------

<p>Serial No.: Exp. Ref./Batch:</p>	<p>0 12313129637</p> 		<p>For use in animals only. Store frozen at ≤ -60°C. Do not vaccinate within 21 days before slaughter. Inactivate unused contents before disposal. See insert for complete information.</p> <p>Pour utilisation chez les animaux seulement. Entreposer au congélateur à une température de -60 °C ou moins. Ne pas vacciner dans les 21 jours précédant l'abattage. Inactiver tout contenu inutilisé avant son élimination. Voir la notice pour l'information complète.</p> <p>Ver las instrucciones completas en el Instructivo adjunto.</p> <p>Manufactured by/Fabriqué par/Fabricado por: Boehringer Ingelheim Animal Health USA Inc. St. Joseph, MO 64506 Tel/Fel.: 1 (888) 637-4251 VLN/PCN 124/10L1.00</p> <p>Distributed in Canada by/Distribué au Canada par/Distribuido en Canada por: Boehringer Ingelheim Animal Health Canada Inc. Boehringer Ingelheim Santé Animale Canada Inc. Burlington ON L7L 5H4</p> 	129623-03
---	--	---	---	-----------

Boehringer Ingelheim Animal Health USA Inc.
3239 Satellite Blvd., Duluth, Georgia 30096-4640
Information 888-325-9167 • Orders 800-325-9167 • Fax 816-236-2717
www.bi-animalhealth.com/swine