

Serial No.:
Exp. Date:

This product has been shown to be effective for the vaccination of healthy swine 2 weeks of age or older against *Salmonella* Choleraesuis and *S. Typhimurium*. Onset of immunity for the *S. Typhimurium* fraction is evident within 14 days of vaccination. The duration of immunity has not been determined. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.

Directions and dosage: Rehydrate the vaccine by adding a portion of the contents of the accompanying diluent to the vaccine vial. Shake well and transfer the reconstituted suspension back into the diluent bottle. Mix with the remaining diluent to complete to a total volume of 200 mL. Shake well. Use entire contents immediately. Administer a single 2 mL dose per pig, either orally or via the drinking water. See insert for complete directions. The need for annual booster vaccinations has not been established for this product; consultation with a veterinarian is recommended.

Precautions: Store at 35-46°F (2-8°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Do not mix with other products, except as specified on this label. Do not use concurrently with antimicrobials effective against *Salmonella* spp. A minimum antibiotic-free period of 3 days pre- and post-vaccination is recommended. All materials used in administering this vaccine must be free of antimicrobial or disinfectant residue to prevent inactivation. In case of anaphylactoid reaction, administer epinephrine. Vaccinated pigs may intermittently shed the vaccine organism in feces for a minimum of 4 weeks after vaccination. This product has not been tested in pregnant animals. In case of human exposure, contact a physician. Inactivate unused contents before disposal.

Boehringer Ingelheim Animal Health USA Inc.
St. Joseph, MO 64506
Phone: 1 (888) 637-4251
VLN/PCN 124/19A2.00

Boehringer Ingelheim

175716-01

Serial No.:
Exp. Date:

Salmonella Choleraesuis-Typhimurium Vaccine
Avirulent Live Culture
ENTERISOL[®] Salmonella T/C FF
For use in swine only
100 Doses/40 mL

This product has been shown to be effective for the vaccination of healthy swine 2 weeks of age or older against *Salmonella* Choleraesuis and *S. Typhimurium*. Onset of immunity for the *S. Typhimurium* fraction is evident within 14 days of vaccination. The duration of immunity has not been established. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.

Directions and dosage: See insert for complete information.

Precautions: Store frozen at ≤ -60 °C. Do not thaw until immediately prior to use. Shake well before use. Do not mix with other products, except as specified on the label. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician. Inactivate unused contents before disposal. Do not use concurrently with antibiotics, including feed antibiotics. A minimum antibiotic-free period of 3 days pre- and post-vaccination is recommended. All materials used in administering this vaccine must be free of antimicrobial or disinfectant residue to prevent inactivation.

Boehringer Ingelheim Animal Health USA Inc.
St. Joseph, MO 64506
Phone: 1 (888) 637-4251
VLN/PCN 124/19A2.01

Boehringer Ingelheim

175801-02

Serial No.:
Exp. Date:

Salmonella Choleraesuis-Typhimurium Vaccine
Avirulent Live Culture
ENTERISOL[®] Salmonella T/C FF
For use in swine only
250 Doses/100 mL

This product has been shown to be effective for the vaccination of healthy swine 2 weeks of age or older against *Salmonella* Choleraesuis and *S. Typhimurium*. Onset of immunity for the *S. Typhimurium* fraction is evident within 14 days of vaccination. The duration of immunity has not been established. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.

Directions and dosage: See insert for complete information.

Precautions: Store frozen at ≤ -60 °C. Do not thaw until immediately prior to use. Shake well before use. Do not mix with other products, except as specified on the label. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician. Inactivate unused contents before disposal. Do not use concurrently with antibiotics, including feed antibiotics. A minimum antibiotic-free period of 3 days pre- and post-vaccination is recommended. All materials used in administering this vaccine must be free of antimicrobial or disinfectant residue to prevent inactivation.

Boehringer Ingelheim Animal Health USA Inc.
St. Joseph, MO 64506
Phone: 1 (888) 637-4251
VLN/PCN 124/19A2.01

Boehringer Ingelheim

175802-02

Serial No.:
Exp. Date:

Salmonella Choleraesuis-Typhimurium Vaccine
Avirulent Live Culture
ENTERISOL[®] Salmonella T/C FF
For use in swine only
500 Doses/200 mL

This product has been shown to be effective for the vaccination of healthy swine 2 weeks of age or older against *Salmonella* Choleraesuis and *S. Typhimurium*. Onset of immunity for the *S. Typhimurium* fraction is evident within 14 days of vaccination. The duration of immunity has not been established. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.

Directions and dosage: See insert for complete information.

Precautions: Store frozen at ≤ -60 °C. Do not thaw until immediately prior to use. Shake well before use. Do not mix with other products, except as specified on the label. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician. Inactivate unused contents before disposal. Do not use concurrently with antibiotics, including feed antibiotics. A minimum antibiotic-free period of 3 days pre- and post-vaccination is recommended. All materials used in administering this vaccine must be free of antimicrobial or disinfectant residue to prevent inactivation.

Boehringer Ingelheim Animal Health USA Inc.
St. Joseph, MO 64506
Phone: 1 (888) 637-4251
VLN/PCN 124/19A2.01

Boehringer Ingelheim

175803-02

Boehringer Ingelheim Animal Health USA Inc.
3239 Satellite Blvd., Duluth, Georgia 30096-4640
Information 888-325-9167 • Orders 800-325-9167 • Fax 816-236-2717
www.bi-animalhealth.com/swine