

This product has been shown to be effective for the vaccination of healthy swine 3 weeks of age or older against porcine circovirus type 2 (PCV2) and *Mycoplasma hyopneumoniae*. The onset of protection against Porcine Circovirus Type 2 is 2 weeks post-vaccination, and the duration of immunity is at least 4 months. The duration of immunity against *M. hyopneumoniae* is at least 26 weeks. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov. The product contains PCV2a, and has been shown to be effective against lymphoid depletion, inflammation and colonization of lymphoid tissue for PCV.

Directions and dosage: Shake each bottle before use. Aseptically mix the full contents of both bottles as per the mixing instructions diagram. See back of this carton. Shake well. Inject a single 2 mL dose intramuscularly. Contact your veterinarian or the manufacturer for information on revaccination frequency.

Porcine Circovirus Vaccine

Type 2, Killed Baculovirus Vector

Mycoplasma Hyopneumoniae Bacterin

FLEXcombo®

250 Doses HSB

Mix to 500 mL

**250
DOSES**



 **Boehringer
Ingelheim**

Precautions: Store at 35 – 46°F (2 – 8°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Do not mix with other products, except as specified on the label. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician.

Preservative: Neomycin.

This package contains one 250 dose (250 mL) vial of Ingelvac CircoFLEX® and one 250 dose (250 mL) vial of Ingelvac MycoFLEX®. The Ingelvac MycoFLEX® is supplied in a headspace bottle (HSB) with extra capacity for the purpose of aseptic mixing.

**Boehringer Ingelheim
Animal Health USA Inc.**
St. Joseph, MO 64506
Phone: 1 (888) 637-4251
VLN/PCN 124/49K5.R1