

FLEXcombo®

Porcine Circovirus Vaccine Type 2, Killed Baculovirus Vector *Mycoplasma Hyopneumoniae* Bacterin

Description

The FLEXcombo® package contains one bottle of Ingelvac CircoFLEX® and one bottle of Ingelvac MycoFLEX®.

FLEXCOMBO is a single-dose porcine circovirus Type 2 (PCV2) and *Mycoplasma hyopneumoniae* vaccine. INGELVAC MYCOFLEX is supplied in a headspace bottle (HSB), with extra capacity for the purpose of convenient mixing with INGELVAC CIRCOFLEX.

Features

- Effective for the vaccination of healthy swine 3 weeks of age or older against porcine circovirus Type 2 (PCV2) and *Mycoplasma hyopneumoniae* (Mhp)
- Single dose, 2 mL
- At least four months duration of immunity for PCV2 and at least 26 weeks duration of immunity for Mhp
- Contains ImpranFLEX®, an effective, aqueous-based (non-oil) polymer adjuvant
- Reduces PCV2 viremia

Benefits

- Low viscosity, highly syringeable
- Highly immunogenic and non-virucidal vaccine that is clear in color, thanks to a state-of-the-art and novel purification process
- Rapid onset of immunity
- Reduced labor costs compared to two-dose vaccine
- Single dose minimizes stress on pigs
- PQA® Plus friendly
- Member of the FLEX Family (nine swine vaccines, multiple combinations); allows you a flexible vaccination scheduling
- NOW, with a new convenient 500-dose presentation (non-HSB)

Dosage & Administration

Shake each bottle before use. Aseptically mix the full contents of both bottles as per the mixing-instruction diagram; see back side of this carton. Shake well. Inject a single 2-mL dose intramuscularly. **Does not need to be tempered.**

Highlights

- Convenient single 2-mL dose
- Rapid onset of immunity as well as long duration of immunity
- Flexible vaccination scheduling

Precautions

Store in dark at 35–46° F (2–8° C).

Do not freeze.

Use entire contents when first opened.

Do not vaccinate within 21 days before slaughter.

Uses

For vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the prevention of lymphoid depletion, inflammation and colonization of lymphoid tissue associated with porcine circovirus Type 2 (PCV2), and as an aid in the prevention of disease caused by *Mycoplasma hyopneumoniae*. The onset of protection against porcine circovirus Type 2 is two weeks post vaccination, and the duration of immunity is at least four months. The duration of immunity against *M. hyopneumoniae* is at least 26 weeks.

Warning

In case of anaphylactic reaction, epinephrine is symptomatic treatment.

Caution: Do not use as diluent for live vaccines.

Product Information

Name	Code	Size
FLEXcombo®	100506	50 dose
	100505	250 dose
	182902	500 dose*

*This presentation does not come with headspace bottle (HSB).



This product has been shown to be effective for the vaccination of healthy swine 3 weeks of age or older against porcine circovirus type 2 (PCV2) and *Mycoplasma hyopneumoniae*. The onset of protection against Porcine Circovirus Type 2 is 2 weeks post-vaccination, and the duration of immunity is at least 4 months. The duration of immunity against *M. hyopneumoniae* is at least 26 weeks. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov. The product contains PCV2a, and has been shown to be effective against lymphoid depletion, inflammation and colonization of lymphoid tissue for PCV.

Directions and dosage: Shake each bottle before use. Aseptically mix the full contents of both bottles as per the mixing instructions diagram. See back of this carton. Shake well. Inject a single 2 mL dose intramuscularly. Contact your veterinarian or the manufacturer for information on revaccination frequency.

Porcine Circovirus Vaccine

Type 2, Killed Baculovirus Vector

Mycoplasma Hyopneumoniae Bacterin

FLEXcombo®

250 Doses HSB

Mix to 500 mL

**250
DOSES**








 **Boehringer
Ingelheim**

Precautions: Store at 35 – 46°F (2 – 8°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Do not mix with other products, except as specified on the label. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician.

Preservative: Neomycin.

This package contains one 250 dose (250 mL) vial of Ingelvac CircoFLEX® and one 250 dose (250 mL) vial of Ingelvac MycoFLEX®. The Ingelvac MycoFLEX® is supplied in a headspace bottle (HSB) with extra capacity for the purpose of aseptic mixing.

**Boehringer Ingelheim
Animal Health USA Inc.**
St. Joseph, MO 64506
Phone: 1 (888) 637-4251
VLN/PCN 124/49K5.R1



SCAN FOR VIDEO
INSTRUCTIONS



ENGLISH



SPANISH

Boehringer Ingelheim Animal Health USA Inc.
3239 Satellite Blvd., Duluth, Georgia 30096-4640
Information 888-637-4251
www.bi-animalhealth.com/swine