BOFHRINGER INGELHEIM

Regulatory Labeling & Graphics - St. loseph, Missouri USA

Type/Size:

PDF:

Revision: 12 Operator: hal

Date:

Art #:

SAP #

Product:

Template #: RA Labeling:

Label/250 ml 291

15Apr20 hal

15Apr20 118702-04 107037-002 BarVac 7/Somnus Dimensions: 3" x 6-63/64" Corner Radius: .125 Master 118701

BLACK PMS 485 PMS 497

PRINTED ON LAMINATE NO VARNISH

Exp. Date:

Indications: This product has been shown to be effective for the vaccination of healthy cattle against Clostridim chauvoei, Cl. septicum, Cl. novyi, Cl. sordellii, Cl. perfringens types C and D, and Haemophilus somnus. This product was licensed prior to the requirement to establish a minimum age for use. The duration of immunity has not been determined. For more information regarding efficacy and safety data, go to productdata.aphis.usda.gov.

Although Cl. perfringens type B is not a significant problem in the U.S.A., immunity may be provided against the beta and epsilon toxins elaborated by Cl. perfringens type B. This immunity is derived from the combination of type C (beta) and type D (epsilon) fractions.

Directions and dosage: Mix well before use. Using aseptic technique, inject 5 mL subcutaneously. Repeat in 21 to 28 days. Contact your veterinarian or the manufacturer for information on revaccination frequency.

Clostridium Chauvoei-Septicum-Novyi-Sordellii-Perfringens Types C & D-Haemophilus Somnus **Bacterin-Toxoid**

BAR-VAC® 7/Somnus



Precautions: Store out of direct sunlight at 35-46°F (2-8°C). Do not freeze. Do not mix with other products. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Transient swelling may occur at the injection site. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician immediately.

Preservative: Phenol

Boehringer Ingelheim Animal Health USA Inc. St. loseph, MO 64506 Phone: 1 (888) 637-4251 VLN/PCN 124/7423.00

