BOEHRINGER INGELHEIM

Regulatory Labeling & Graphics - St. Joseph, Missouri USA

Revision: 9
Operator: ars
Date: 18Jun19
Art #: 120302-05
SAP #: 106168-002

Type/Size: Label/250 mL Template #: 291 Dimensions: 3" x 6-63/64"

 Corner Radius:
 .125

 Master:
 120301

 PDF:
 18Jun19 ars

PMS 485

PRINTED ON LAMINATE. NO VARNISH.

Bar-Vac CD/T

Exp. Date

Product:

Indications: This product has been shown to be effective for the vaccination of healthy sheep, goats and cattle against Clostridium perfringens types C and D and Cl. tetani. This product was licensed prior to the requirement to establish a minimum age for use. The duration of immunity has not been determined. For more information regarding efficacy and safety data, go to productdata.aphis. usda.gov.

Although *Cl. perfringens* type B is not a significant problem in the U.S.A., immunity may be provided against the beta and epsilon toxins elaborated by *Cl. Perfringens* type B. This immunity is derived from the combination of type C (beta) and type D (epsilon) fractions.

Directions and dosage: Mix well before use. Cattle: Using aseptic technique, inject 5 mL subcutaneously. Repeat in 21 to 28 days. Sheep and Goats: Using aseptic technique, inject 2 mL subcutaneously. Repeat in 21 to 28 days.




Contact your veterinarian or the manufacturer for information on revaccination frequency.

Precautions: Store out of direct sunlight at 35-46°F (2-8°C). Do not freeze. Do not mix with other products. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Transient swelling may occur at the injection site. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician immediately.

Preservative: Phenol.

Boehringer Ingelheim Animal Health USA Inc. St. Joseph, MO 64506 Phone: 1 (888) 637-4251 VLN/PCN 124/8304.00



120302-05

