

BOEHRINGER INGELHEIM

Regulatory Labeling & Graphics - St. Joseph, Missouri USA

Revision:	10	Type/Size:	Label/250 mL
Operator:	ars	Template #:	291
Date:	05Sep19	Dimensions:	3" x 6-63/64"
Art #:	110902-04	Corner Radius:	.125
SAP #:	106157-002		
Product:	Bar-Vac CD	PDF:	05Sep19 ars

BLACK

PMS 485

PMS 652

PRINTED ON LAMINATE. NO VARNISH.

Serial No.:
Exp. Date:

Indications: This product has been shown to be effective for the vaccination of healthy sheep and cattle, including pregnant and lactating cows, against *Clostridium perfringens* types C and D. This product was licensed prior to the requirement to establish a minimum age for use. The duration of immunity has not been determined. For more information regarding efficacy and safety data, go to productdata.aphis.usda.gov.

Although *Cl. perfringens* type B is not a significant problem in the U.S.A., immunity may be provided against the beta and epsilon toxins elaborated by *Cl. Perfringens* type B. This immunity is derived from the combination of type C (beta) and type D (epsilon) fractions.

Directions and dosage: Mix well before use. **Sheep:** Using aseptic technique, inject 2 mL subcutaneously or intramuscularly. Repeat in 21 to 28 days. **Cattle:** Using aseptic technique, inject



Clostridium Perfringens Types C & D Toxoid

BAR-VAC® CD250 mL/50 Cattle Doses
or 125 Sheep Doses

5 mL subcutaneously or intramuscularly. Repeat in 21 to 28 days.

Contact your veterinarian or the manufacturer for information on revaccination frequency.

Precautions: Store out of direct sunlight at 35-46°F (2-8°C). Do not freeze. Do not mix with other products. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Transient swelling may occur at the injection site. In case of anaphylactoid reaction, administer epinephrine. In case of human exposure, contact a physician.

Preservative: Phenol
Boehringer Ingelheim Animal Health USA Inc.
St. Joseph, MO 64506
Phone: 1 (888) 637-4251
VLN/PCN 124/8201.01



110902-04