Boehringer Ingelheim Regulatory Labeling & Graphics - St. Joseph, Missouri USA

Revision: 11 Type/Size: Label/10 Dose/50 mL

Cl. septicum, Cl. haemolyticum, Cl. novyi,

This product was licensed prior to the

Although CL nerfringens type B is not a

combination of type C (beta) and

Cattle: Using aseptic technique, inject

aphis usda gov.

This immunity is derived from the

N

type D (epsilon) fractions.

hal Template #: 299 Operator: 19lun19 Dimensions: 1-1/2" x 4"

Date: 124201-03 Corner Radius: Art #: .125

106210-002 MASTER

Product: Bar Vac 8 PDF: 19lun19 hal


No varnish, printed on laminate,

Date

0

SAP #:

BLACK PMS 485 PMS 690



subcutaneously. Repeat in 21 to 28 days. Contact your veterinarian or the manufacturer for information on revaccination frequency.

Precautions: Store out of direct sunlight at 35-46°F (2-8°C). Do not freeze. Do not mix with other products. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Transient swelling may occur at the injection site. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician.

Preservative: Phenol

Boehringer Ingelheim Animal Health USA, Inc. St. Joseph, MO 64506 Phone: 1 (888) 637-4251 VI N/PCN 124/7160.00

Boehringer Ingelheim Boehringer 124201-03