

# BOEHRINGER INGELHEIM

Regulatory Labeling & Graphics - St. Joseph, Missouri USA

Revision:	13	Type/Size:	Label/50 dose
Operator:	hal		
RA Labeling:		Template #:	269
Date:	22Jan20	Dimensions:	1.77" x 5"
Art #:	133302-02	Corner Radius:	.125
SAP #:	106376-002	MASTER:	133301
Product:	Caliber 7	PDF:	22Jan20 hal

BLACK

PMS 485

PMS 357


PRINTED ON LAMINATE. NO VARNISH.

Serial  
No.:  
Exp.  
Date:

**Indications:** This product has been shown to be effective for the vaccination of healthy sheep and cattle, including pregnant and lactating cows, against *Clostridium chauvoei*, *Cl. septicum*, *Cl. novyi*, *Cl. sordellii*, and *Cl. perfringens* types C and D. This product was licensed prior to the requirement to establish a minimum age for use. The duration of immunity has not been determined. For more information regarding efficacy and safety data, go to [productdata.aphis.usda.gov](http://productdata.aphis.usda.gov).

Although *Cl. perfringens* type B is not a significant problem in the U.S.A., immunity may be provided against the beta and epsilon toxins elaborated by *Cl. perfringens* type B. This immunity is derived from the combination of type C (beta) and type D (epsilon) fractions.

**Directions and dosage:** Mix well before use. Using aseptic technique, inject 2 mL



subcutaneously. Repeat in 21 to 28 days. Contact your veterinarian or the manufacturer for information on revaccination frequency.

**Precautions:** Store out of direct sunlight at 35-46°F (2-8°C). Do not freeze. Do not mix with other products. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant sheep. In case of human exposure, contact a physician immediately.

Boehringer Ingelheim Animal Health USA Inc.  
St. Joseph, MO 64506  
Phone: 1 (888) 637-4251  
VLN/PCN 124/7410.02



Boehringer  
Ingelheim

133302-02