

BOEHRINGER INGELHEIM

Regulatory Labeling & Graphics - St. Joseph, Missouri USA

Revision:	17	Type/Size:	Label/50 dose
Operator:	hal		
RA Labeling:		Template #:	269
Date:	10Aug20	Dimensions:	1.77" x 5"
Art #:	118402-04	Corner Radius:	.125
SAP #:	107029-002		
Product:	Citadel L5 HB		

BLACK

PMS 258

PMS 202

PRINTED ON LAMINATE. NO VARNISH.


Serial No.:
Exp. Date:

0
1231311843
3

Indications: This product has been shown to be effective for the vaccination of healthy cattle or swine against *Leptospira canicola*, *L. grippityphosa*, *L. hardjo*, *L. icterohaemorrhagiae*, and *L. pomona*. This product was licensed prior to the requirement to establish a minimum age for use. The duration of immunity has not been determined. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.

This product has also been shown to be effective against urinary shedding of *L. borgpetersenii* serovar *hardjo* (type *hardjo-bovis*). Vaccinated heifers subsequently exposed to *L. borgpetersenii* serovar *hardjo* (type *hardjo-bovis*) have been shown to clear renal infections within 8 weeks of exposure.

Directions and dosage: Mix well before use. Using aseptic technique, inject 2 mL subcutaneously or intramuscularly in healthy cattle, or intramuscularly in swine. A second



dose is recommended 3 to 6 weeks later. Contact your veterinarian or the manufacturer for information on revaccination frequency.

Precautions: Store out of direct sunlight at 35-46°F (2-8°C). Do not freeze. Do not mix with other products. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician.

Preservatives: Neomycin and thimerosal.

Boehringer Ingelheim Animal Health USA Inc.
St. Joseph, MO 64506
Phone: 1 (888) 637-4251
VLN/PCN 124/2665.00



118402-04