BOEHRINGER INGELHEIM

Regulatory Labeling & Graphics - St. Joseph, Missouri USA

Revision: 12

SAP #:

Type/Size: Label/50 dose

hal Operator: RA Labeling: 10Aug20 Date: Art #: 129102-02

Template #: 269 Dimensions: 1.77" x 5" Corner Radius: .125

MASTER 129101


Citadel VI 5 HB Product:

107189-002

PRINTED ON LAMINATE. NO VARNISH.

BLACK PMS 258 PMS 072

Serial No.: Exp. Date: Indications: This product has been shown to be effective for the vaccination of healthy cattle against Campylobacter fetus var. venerealis, Leptospira canicola, L. grippotyphosa, L. hardjo, L. icterohaemorrhagiae, and L. pomona. This product was licensed prior to the requirement to establish a minimum age for use. The duration of immunity has not been determined. For more information regarding efficacy and safety data go to productdata. aphis.usda.gov. This product has also been shown to be effective against urinary shedding of L. borqpetersenii serovar hardjo (type hardjo-bovis) and against infertility, delayed conception, or abortion due to C. fetus var. venerealis. Vaccinated animals subsequently exposed to L. borapetersenii serovar hardio (type hardio-hovis) have been shown to clear renal infections within 8 weeks of \sim Directions and dosage; Mix well before use.



Using aseptic technique, inject 2 mL subcutaneously. Repeat in 14 to 28 days. Contact your veterinarian or the manufacturer for information on revaccination frequency.

Precautions: Store out of direct sunlight at 35-46°F (2-8°C). Do not freeze. Do not mix with other products. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction. administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician.

Preservatives: Neomycin and thimerosal.

Boehringer Ingelheim Animal Health USA, Inc. St. Joseph, MO 64506 Phone: 1 (888) 637-4251 VLN/PCN 124/2863.01

Boehringer Ingelheim Boehringer 129102-02