

 Boehringer Ingelheim GO STJ	<b>Code: 126705-07</b> SAP #: 107089-004
INITIALS: HL	Version: 1      28Mar22
<b>Product</b> Component Type Size:	<b>EXPRESS FP 5</b> Carton 50 Doses/100 mL
Country(s):	US
Dimensions Flat:	3.938" x 2.063" x 4.250"
Dimensions Folded:	NA
Template #:	842
Corner Radius:	NA
<b>DIELINE:</b>	L071796E_C
RA Labeling:	
Printer Notes:	

COLORS
BLACK
BLACK-70%
PMS 135
PMS 072

126705-07

Serial No:  
Exp. Date:50 Doses/Rehydrate with 100 mL  
**EXPRESS® FP 5**

**Indications:** This product has been shown to be effective for the vaccination of healthy cattle, including pregnant cows, against reproductive and respiratory disease due to infectious bovine rhinotracheitis (IBR) and bovine virus diarrhea types 1 and 2 (BVD1 & BVD2), and against respiratory disease due to parainfluenza 3 (PI3) and bovine respiratory syncytial virus (BRSV). This product was licensed prior to the requirement to establish a minimum age for use. The duration of immunity against IBR-induced abortion and against BVD1b & BVD2, including persistently infected calves, is at least 12 months. The duration for the remaining organisms has not been determined. For more information regarding efficacy and safety data go to [productdata.aphis.usda.gov](http://productdata.aphis.usda.gov).

The product has also been shown to be effective against persistently infected calves due to BVD1b & BVD2. This vaccine may be used in pregnant cows or calves nursing pregnant cows, provided the cows were vaccinated pre-breeding, within the past 12 months, according to label instructions with any Express FP vaccine.

**Composition:** The amber glass vial contains IBR, BVD1 (Singer 1a cytopathic), BVD2 (296 cytopathic), PI3, and BRSV modified live viruses (MLV). The plastic vial contains an adjuvant system. Neomycin and thimerosal added as preservatives.

**Directions and dosage:** Mix the accompanying bottle of diluent. Rehydrate the vaccine by aseptically adding the diluent to the vaccine vial. Mix the rehydrated vaccine and use immediately. Using aseptic technique, inject 2 mL subcutaneously or intramuscularly. If using the subcutaneous route, inject in front of the shoulder and midway of the neck, away from the suprascapular lymph node. For cows and heifers, using aseptic technique, inject a single 2 mL dose subcutaneously or intramuscularly at or about 4 weeks prior to breeding. If initial vaccination, repeat with any Express® vaccine containing BRSV MLV in 14 to 28 days. Contact your veterinarian or the manufacturer for information on revaccination frequency.

**Precautions:** Store out of direct sunlight at 35–46°F (2–8°C). Do not freeze. Do not mix with other products, except as specified on this label. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Injection site swelling may occur. Stressed cattle should not be vaccinated. In case of anaphylactoid reaction, administer epinephrine. In case of human exposure, contact a physician. Inactivate unused contents before disposal.

Fetal health risks associated with vaccination of pregnant animals with this vaccine cannot be unequivocally determined by clinical trials conducted for licensure. Appropriate strategies to address the risks associated with vaccine use in pregnant animals should be discussed with a veterinarian. No vaccine can be expected to have 100% efficacy under all conditions. A small number of calves persistently infected with BVD may have a devastating effect on herd health.

**Note:** It is possible that healthy-appearing cattle can be persistently infected with or incubating virulent BVD virus at the time of vaccination.

## Bovine Rhinotracheitis- Virus Diarrhea- Parainfluenza 3- Respiratory Syncytial Virus Vaccine

Modified Live Virus

**EXPRESS® FP 5**

50 Doses/Rehydrate with 100 mL

50  
DOSES
 **Boehringer  
Ingelheim**

In view of these findings and suggested causes, BVD vaccine is contraindicated in persistently infected cattle and use should be limited only to healthy, immunocompetent, unstressed cattle.

This package contains one 50 dose vial of MLV vaccine and one 100 mL vial of adjuvanted diluent.

**Boehringer Ingelheim  
Animal Health USA Inc.**  
St. Joseph, MO 64506  
Phone: 1 (888) 637-4251  
VLN/PCN 124/1181.24



Express® is a registered trademark of Boehringer Ingelheim  
Animal Health USA Inc.



126705-07

GLUE