 Boehringer Ingelheim	Code: 124600-01
GO STJ	SAP #: 165124-002
INITIALS: KS-HL	Version: 1 01Apr21
Product	FLEXParvoPRRS
Component Type Size:	Carton 50ds
Country(s):	US
Dimensions Flat:	3.938" x 2.063" x 4.250"
Dimensions Folded:	NA
Template #:	842
Corner Radius:	NA
DIELINE:	L071796E_C
RA Labeling:	
Printer Notes:	NA

COLORS	
BLACK	PMS 499
PMS 10147	PMS 354
PMS 287	
PMS 877	

10-009721




Serial No.:
Exp. Date:

FLEXParvoPRRS®
50 Doses/Rehydrate to 100 mL

This product has been shown to be effective for the vaccination of healthy swine 6 months of age or older against the reproductive failure caused by porcine parvovirus (PPV) and reproductive disease caused by Porcine Reproductive and Respiratory Syndrome (PRRS) virus. The duration of immunity is at least 4 months for the reproductive form of PRRS virus. The duration of immunity has not been determined for porcine parvovirus. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.

Directions and dosage: Rehydrate the vaccine by adding a portion of the contents of the ReproCyc® ParvoFLEX vial to the Ingelvac PRRSFLEX® MLV vaccine vial. Shake well and transfer the reconstituted suspension back into the ReproCyc® ParvoFLEX vial. Mix with the remaining ReproCyc® ParvoFLEX contents to complete the total volume. Shake well and use immediately. Using aseptic technique, inject a single 2 mL dose of vaccine intramuscularly. Administer a second 2 mL intramuscular dose of ReproCyc® ParvoFLEX 3 weeks later. For more information on revaccination frequency, consult your veterinarian.



Precautions: Store out of direct sunlight at 35 - 46°F (2 - 8°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Do not mix with other products, except as specified on this carton. In case of anaphylactoid reaction, administer epinephrine. For use in PRRS virus positive herds only. Do not vaccinate breeding age boars. PRRS vaccine virus may be shed and transmitted to other populations of swine which are in contact with vaccinated swine. The duration of potential virus transmission may vary. Use of the vaccine in herds intended to remain PRRS virus seronegative is contraindicated. Introduction of vaccinated pigs into herds intended to remain



Porcine Reproductive & Respiratory Syndrome-Parvovirus Vaccine

Reproductive Form, Modified Live Virus, Killed Baculovirus Vector

FLEXParvoPRRS®

50 Doses/Rehydrate to 100 mL



PRRS virus seronegative is contraindicated. Vaccine is safe in pregnant swine. Specific herds may be particularly sensitive. Fetal health risks associated with the vaccination of pregnant animals with this vaccine cannot be unequivocally determined during clinical trials conducted for licensure. Appropriate strategies to address the risks associated with vaccine use in pregnant animals should be discussed with a veterinarian. In case of human exposure, contact a physician. Inactivate unused contents before disposal.

Preservative: Neomycin

Boehringer Ingelheim Animal Health USA Inc.
St. Joseph, MO 64506
1 (888) 637-4251
VLN/PCN 124/1P29.R0



124600-01