

Ingelvac CircoFLEX®

Porcine Circovirus Vaccine Type 2, Killed Baculovirus Vector

Description

Ingelvac CircoFLEX® is a single-dose porcine circovirus Type 2 (PCV2) vaccine.

Features

- Single dose, 1 mL
- Onset of immunity within two weeks of vaccination
- At least four months' duration of immunity
- Reduces PCV2 viremia
- Contains ImpranFLEX®, an effective, aqueous-based (non-oil) polymer adjuvant

Benefits

- PQA® Plus friendly
- Member of the FLEX Family (nine swine vaccines, multiple combinations); allows you a flexible vaccination scheduling
- Highly immunogenic and non-virucidal vaccine that is clear in color, thanks to a state-of-the-art and novel purification process
- NOW with a new convenient 500-dose presentation

Dosage & Administration

Shake well before use. Using aseptic technique, administer a single 1-mL dose by intramuscular injection. **Does not need to be tempered.**

Precautions

Store in dark at 35–46° F (2–8° C).

Do not freeze.

Use entire contents when first opened.

Do not vaccinate within 21 days before slaughter.

Uses

For vaccination of healthy, susceptible pigs 3 weeks of age or older as an aid in the prevention of lymphoid depletion, inflammation and colonization of lymphoid tissue, and as an aid in reduction of the magnitude of viremia associated with porcine circovirus Type 2 (PCV2).

Warning

In case of anaphylactic reaction, epinephrine is symptomatic treatment.

Product Information

Name	Code	Size
Ingelvac	138708	50 dose
CircoFLEX®	138518	100 dose
	138710	250 dose
	131167	500 dose

Highlights

- Convenient single 1-mL dose
- At least four months DOI
- Highly immunogenic and non-virucidal vaccine
- Flexible vaccination scheduling




Serial No.:
Exp. Date:

This product has been shown to be effective for the vaccination of healthy swine 3 weeks of age or older against porcine circovirus type 2 (PCV2). The onset of protection is 2 weeks post-vaccination. The duration of immunity is at least 4 months. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.

The product contains PCV2a, and has been shown to be effective against lymphoid depletion, inflammation and colonization of lymphoid tissue, and the magnitude of viremia.

Directions and dosage: Shake well before use. Using aseptic technique, administer a single **1 mL** dose by intramuscular injection.



Precautions: Store out of direct sunlight at 35 - 46°F (2 - 8°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Do not mix with other products. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician.

Preservative: Neomycin

**Boehringer Ingelheim
Animal Health USA Inc.**
St. Joseph, MO 64506
Phone: 1 (888) 637-4251
VLN/PCN 124/19K5.R0



130904-07

Boehringer Ingelheim Animal Health USA Inc.
3239 Satellite Blvd., Duluth, Georgia 30096-4640
Information 888-637-4251
www.bi-animalhealth.com/swine