



GO STI  
JR - HL

Code: 160802-04  
SAP #: 107302-003

Version + Date: 17 - 23Feb21 MASTER: 160801

Product:	Triangle 5
Component Type/Size:	Label/50ds/100mL
Country(s):	US
Dimensions FLAT:	1.77" x 5.000"
Dimensions FOLDED:	N/A
Template #:	269
Dieline:	N/A
Corner Radius:	.125"

RA Labeling:

Printer Notes:

Printed on Laminate, no varnish

Colors

PMS 362

BLACK

PMS 205


Serial No.:

Exp. Date:

0  
1  
2  
3  
1  
3  
1  
1  
6  
0  
8  
3  
8

**Indications:** This product has been shown to be effective for the vaccination of healthy cattle, including pregnant cows, against infectious bovine rhinotracheitis (IBR), bovine virus diarrhea types 1 and 2 (BVD1 & BVD2), parainfluenza 3 (PI3) and bovine respiratory syncytial virus (BRSV). This product was licensed prior to the requirement to establish a minimum age for use. The duration of immunity has not been determined. For more information regarding efficacy and safety data, go to [productdata.aphis.usda.gov](http://productdata.aphis.usda.gov).

**Directions and dosage:** Mix well. Inject one 2 mL dose intramuscularly or subcutaneously, using aseptic technique. Repeat in 14 to 28 days. Protect animals from exposure for at least 14 days after vaccination. Contact your veterinarian



or the manufacturer for information on revaccination frequency.

**Precautions:** Store out of direct sunlight at 35–46°F (2–8°C). Do not freeze. Do not mix with other products. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Transient swelling may occur at the injection site. In case of anaphylactoid reaction, administer epinephrine. In case of human exposure, contact a physician.

**Preservatives:** Gentamicin, neomycin, polymyxin B, and thimerosal.

**Boehringer Ingelheim  
Animal Health USA Inc.**  
St. Joseph, MO 64506  
Phone: 1 (888) 637-4251  
VLN/PCN 124/1185.23

160802-04