

more life to love

VETMEDIN® (pimobendan) Chewable Tablets should be administered orally at a total daily dose of 0.5 mg/kg of body weight.

Hours

The total daily dose might be divided into two portions that are not necessarily equal, and the portions should be given approximately 12 hours apart

Any suitable combination of whole- and/or half-tablets may be used

Dosage chart					
Tablet size*	Body weight	Total daily dosage	Bottle duration	Morning	Evening
1.25 mg	2.5 kg/5.5 lb	1.25 mg	50 days		
	3.75 kg/8.3 lb	1.875 mg	33 days		
2.5 mg	5 kg/11 lb	2.5 mg	50 days		
	7.5 kg/16.5 lb	3.75 mg	33 days		
5 mg	10 kg/22 lb	5 mg	50 days		
	15 kg/33 lb	7.5 mg	33 days		
	20 kg/44 lb	10 mg	25 days		
10 mg	30 kg/66 lb	15 mg	33 days		
	40 kg/88 lb	20 mg	25 days		

Please note: These examples given in the dosage chart do not necessarily depict the only way to provide the correct dose. Other combinations of whole- and half-tablets may also be appropriate.

For more information, contact your Boehringer Ingelheim sales representative.

IMPORTANT SAFETY INFORMATION: VETMEDIN® (pimobendan) is for use in dogs with clinical evidence of heart failure only. The most common side effects reported in field studies were poor appetite, lethargy, diarrhea, dyspnea, azotemia, weakness, and ataxia. If side effects should occur, pet owners should contact their veterinarian. For complete product information, see the full prescribing information or visit www.VETMEDINClinic.com



VETMEDIN® is a registered trademark of Boehringer Ingelheim Vetmedica GmbH, used under license. ©2024 Boehringer Ingelheim Animal Health USA Inc., Duluth, GA. All rights reserved. US-PET-1074-2024.

