

PCV2d

Porcine Circovirus Vaccine Type 2, Killed Baculovirus Vector

Description

d-FENSE is a single-dose Porcine Circovirus Type 2 (PCV2) vaccine.

Advantages

- Onset of immunity within two weeks of vaccination
- At least four months duration of immunity
- Single dose, 1 mL
- Low risk of adverse reactions and stress on pigs due to one-dose administration
- PQA® Plus friendly
- Contains ImpranFLEX®, an effective, aqueous-based (non-oil) polymer adjuvant
- Flexible vaccination scheduling
- Reduces PCV2 viremia.

Benefits

d-FENSE is a single-dose PCV2 vaccine that can be administered to pigs 3 weeks of age and older while providing a DOI of at least four months.

Dosage & Administration

Shake well before use. Using aseptic technique, administer a single 1-mL dose by intramuscular injection.

Precautions

Store in dark at 35–46° F (2–8° C). Do not freeze. Use entire contents when first opened.

Uses

For vaccination of healthy, susceptible pigs 3 weeks of age or older as an aid in the prevention of lymphoid depletion, inflammation and colonization of lymphoid tissue, and as an aid in reduction of the magnitude of viremia associated with Porcine Circovirus Type 2 (PCV2). The onset of protection is two weeks post vaccination. The duration of immunity is at least four months.

Warning


Do not vaccinate within 21 days before slaughter. In case of anaphylactic reaction, epinephrine is symptomatic treatment

Highlights

- Convenient single 1-mL dose
- For pigs as young as 3 weeks of age
- At least four months DOI
- Low risk of adverse reactions

Product Information

Name	Code	Size
Porcine Circovirus Vaccine Type 2, Killed Baculovirus Vector	162664	100mL




Serial No.:
Exp. Date:

Composition: This product contains PCV2d.

This product has been shown to be effective for the vaccination of healthy pigs 3 weeks of age or older against porcine circovirus type 2d (PCV2d). The onset of immunity is 28 days. The duration of immunity is not determined. For more information regarding efficacy and safety data, go to productdata.aphis.usda.gov.

The efficacy of the product was demonstrated based on lymphoid depletion, lymphoid colonization, and viremia, when administered intramuscularly as a single 1 mL dose to 3-week old pigs.

Directions and dosage: Shake well before use. Using aseptic technique, administer a single 1 mL dose by intramuscular injection. The need for annual booster vaccinations has not been established for this product; consultation with a veterinarian is recommended.



Precautions: Store at 35-46°F (2-8°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Do not mix with other products. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician.

Preservative: Neomycin

Boehringer Ingelheim Animal Health USA Inc.
St. Joseph, MO 64506
Phone: 1 (888) 637-4251
VLN/PCN 124/19K5.R4



**Boehringer
Ingelheim**

131001-01

Boehringer Ingelheim Animal Health USA Inc.
3239 Satellite Blvd., Duluth, Georgia 30096-4640
Information 888-637-4251
www.bi-animalhealth.com

US-POR-0297-2021

IMPRANFLEX® is a registered trademark of Boehringer Ingelheim Vetmedica GmbH, used under license.
©2022 Boehringer Ingelheim Animal Health USA Inc., Duluth, GA. All Rights Reserved.