



|   |                           |                |
|---|---------------------------|----------------|
|  <b>Boehringer<br/>Ingelheim</b> | <b>Code: 176026-01</b>    |                |
| <b>GO STJ</b>   | SAP #: 180862-002         |                |
| <b>INITIALS: KS-HL</b>  | <b>Version: 1</b>         | <b>13Dec21</b> |
| <b>Product</b>  | <b>ReproCyc ParvoFLEX</b> |                |
| Component Type Size:  | Label 50ds / 100 mL       |                |
| Country(s):   | US                        |                |
| Dimensions Flat:  | 1.766" x 5.000"           |                |
| Dimensions Folded:  | NA                        |                |
| Template #:   | 269                       |                |
| Corner Radius:  | .125"                     |                |
| <b>DIELINE:</b>   | NA                        |                |
| <b>RA Labeling:</b>   |                           |                |
| <b>Printer Notes:</b>   | NA                        |                |

| COLORS           |                |
|------------------|----------------|
| <b>BLACK</b>     | <b>PMS 354</b> |
| <b>PMS 10147</b> | <b>PMS 287</b> |
| <b>PMS 499</b>   |                |

Serial No.:  
Exp. Date:

This product has been shown to be effective for the vaccination of healthy sows and gilts 6 months of age or older against reproductive failure caused by porcine parvovirus (PPV). The duration of immunity has not been determined. For more information regarding efficacy and safety data go to [productdata.aphis.usda.gov](http://productdata.aphis.usda.gov).

**Directions and dosage:** Shake well before use. Using aseptic technique, administer a single 2 mL dose by intramuscular injection. Administer a second intramuscular dose 3 weeks later. For more information on revaccination frequency, consult your veterinarian. Safe for use in pregnant animals.



**Precautions:** Store at 35-45°F (2-8°C). Do not freeze. Do not mix with other products. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction, administer epinephrine. In case of human exposure, contact a physician.

**Preservative:** Neomycin

**Boehringer Ingelheim  
Animal Health USA Inc.**  
St. Joseph, MO 64506  
1 (888) 637-4251  
VLN/PCN 124/18P5.R0

 **Boehringer  
Ingelheim**

176026-01